

2

Chlamydia

Patient name: _____ Admission: _____

NRS
DATE INITIAL

--	--

- I. The client/caregiver can define chlamydia.
 - A. It is a sexually transmitted disease caused by the bacterium *Chlamydia trachomatis*.
 - B. It can infect both men and women.
 - C. It can damage female reproductive organs, including cause infertility. Men rarely have complications.
 - D. The bacteria infect the cervix and urethra and can spread to fallopian tubes.
 - E. Chlamydia can infect the rectum. Even mouth and throat infection can occur if having oral sex with infected partner.
 - F. Testing for chlamydia is done by obtaining cultures of cervix or penis.
- II. The client/caregiver can list symptoms of chlamydia.
 - A. This is a “silent” disease because the majority of infected people have no symptoms.
 - B. If symptoms appear, they will show within 1 to 3 weeks after exposure.
 - C. Symptoms for women
 - Abnormal vaginal discharge
 - Burning sensation when urinating
 - Lower abdominal pain
 - Low back pain
 - Nausea
 - Fever
 - Pain during intercourse
 - Bleeding between menstrual periods
 - D. Symptoms for men
 - Discharge from penis
 - Burning sensation when urinating
 - Burning or itching around opening of the penis
 - E. Even mouth and throat infection can occur if having oral sex with infected partner.
 - F. If rectal infection, rectal pain, discharge, or bleeding may occur.
- III. The client/caregiver can list treatment options.
 - A. Antibiotic therapy is administered.
 - B. Sex partners of infected person should be evaluated, tested, and treated.

NRS
DATE INITIAL

--	--

- C. Infected persons should abstain from sexual intercourse until they and their sex partners have completed treatment.
- D. Retesting can be considered 3 to 4 months after treatment.
- IV. The client/caregiver can describe preventive measures.
 - A. Abstain from sexual contact.
 - B. Be in a long-term mutually monogamous relationship.
 - C. Have correct and consistent use of latex male condoms.
 - D. Have annual screening for sexually active women 25 years of age or younger.
 - E. Have annual screening for women with new sex partner or multiple sex partners.
 - F. All pregnant women should be screened.

RESOURCES

Community Health Clinic

Healthy People 2010

www.health.gov/healthypeople

Centers for Disease Control and Prevention

www.cdc.gov/std/

REFERENCES

- Ackley, B. J., & Ladwig, G. B. (2006). *Nursing diagnosis handbook: A guide to planning care*. St. Louis: Mosby Inc.
- Centers for Disease Control and Prevention. (2002). Sexually transmitted diseases treatment guidelines. MMWR 2002;51 (no. RR-6).
- Centers for Disease Control and Prevention. (2005, September). Sexually transmitted disease surveillance. Atlanta, GA: U.S. Department of Health and Human Services.
- Cohen, B. J., & Taylor, J. J. (2005). *Memmler's the human body in health and disease* (10th ed.). Philadelphia: Lippincott Williams & Wilkins.
- Hunt, R. (2005). *Introduction to community based nursing*. Philadelphia: Lippincott Williams & Wilkins.
- Perry, A., & Potter, P. (2006). *Clinical nursing skills & technique*. St. Louis: Mosby Inc.
- Timby, B. K., & Smith, N. C. (2003). *Introductory medical-surgical nursing* (8th ed.). Philadelphia: J. B. Lippincott Williams & Wilkins.