

**2**

# Anticoagulant and Antiplatelet Agents

Patient name: \_\_\_\_\_

NRS  
DATE INITIAL

--	--

Admission: \_\_\_\_\_

NRS  
DATE INITIAL

--	--

- I. The client/caregiver can define the purpose and action of oral anticoagulant and antiplatelet agents.
  - A. The anticoagulant and antiplatelet medication is used to treat or prevent a blood clot that could result in a heart attack, stroke, and other life-threatening problems.
  - B. The medications discussed here are to be taken as oral medication.
  
- II. The client/caregiver can list important assessments and evaluations used when taking oral anticoagulants.
  - A. Report any signs of bleeding (nosebleeds, bleeding gums, or blood in urine) or bruising promptly.
  - B. Avoid taking drugs that may increase the risk of bleeding, such as aspirin or nonsteroidal antiinflammatory drugs without approval of physician.
  - C. Instruct client to avoid activities with high risk of injury.

- D. Male clients should use electric razor.
- E. Take medication exactly as prescribed.
- F. Have blood work done at times specified by physician.
- G. Keep all follow-up visits with physician.
- H. Keep record of any medication changes your physician may give over the phone as a result of recent laboratory tests.
- I. Read food labels. Any foods or supplements that contain vitamin K may impair the effectiveness of anticoagulant medication.

## REFERENCES

- Ackley, B. J., & Ladwig, G. B. (2006). *Nursing diagnosis handbook: A guide to planning care*. St. Louis: Mosby Elsevier.
- Deglin, J. H., & Vallerand, A. H. (2001). *Davis's drug guide for nurses*. Philadelphia: F.A. Davis Company.
- Nursing 2006 drug handbook*. (2006). Philadelphia: Lippincott Williams and Wilkins.
- Rice, J. (1998). *Medications mathematics for the nurse*. Albany, NY: Delmar.